

PALM INTRAXIET

Day: Wednesday Date: 10/4/2006 Time: 13:00:46

Inventor Information for 60/447558

Inventor Name	n/	City		State/Country	
MARQUIS, ROBERT W.	1 N	COLLEGI	EVILLE	PENNSYLVANIA	
Appin Info Contents P	etition Info	Atty/Agent I	info Continuity/	Reexam Foreign Data	Invent
Search Another: Applicati	on#	Search	or Patent#	Search	
PCT /	/	Search	or PG PUBS#	Search	
Attorney D	ocket#		Search		,
Bar Code #		Sear	ch		

To go back use Back button on your browser toolbar.

Back to PALM | ASSIGNMENT | OASIS | Home page

FILE 'CAPLUS' ENTERED AT 14:10:43 ON 04 OCT 2006 L1	·
L2 0 S US10/772817/PN L3 0 S MARQUIS AND QUINOLINE-6-CARBOXYLIC	
L3 0 S MARQUIS AND QUINOLINE-6-CARBOXYLIC	
L3 0 S MARQUIS AND QUINOLINE-6-CARBOXYLIC	
L4 95 S QUINOLINE-6-CARBOXYLIC	
L5 1 S L4 AND AZEPAN-4-YL	
L6 1 S US 2004-192674/PN	
L7 4 S 350796-38-2/RN OR 350796-41-7/RN OR 764650-55-7/RN	
L8 4 DUP REM L7 (0 DUPLICATES REMOVED)	

ANSWER 1 OF 1 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2004:803931 CAPLUS

DOCUMENT NUMBER: 141:295878

TITLE: Preparation of aminoazepanones as Cathepsin L

inhibitors

INVENTOR(S): Marquis, Robert W.

PATENT ASSIGNEE(S): USA

SOURCE: U.S. Pat. Appl. Publ., 13 pp.

CODEN: USXXCO

DOCUMENT TYPE:

Patent

Ι

LANGUAGE:

GΙ

English

FAMILY ACC. NUM. COUNT:

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2004192674 PRIORITY APPLN. INFO.:	A1	20040930	US 2004-772817 US 2003-447558P P	20040205 < 20030214
OTHER SOURCE(S):	MARPAT	141:295878		

 AB The title compds. I [R1 = substituted aminoalkylcarbonyl; R2 = H, alkyl, arylalkyl, etc.; R3 = H, alkyl, cycloalkyl, arylalkyl, etc.; R4 = H, alkyl, cycloalkyl, etc.] and their pharmaceutically acceptable salts are prepared and disclosed as inhibitors of Cathepsin L. Thus, e.g., II was prepared in a multistep synthesis employing N-Boc-phenylalanine. Consequently they are useful for preventing or treating diseases in which cathepsin L is implicated, such as rheumatoid arthritis or inhibition of pos. selection of CD4 + T-cells by cortical thymic epithelial cells.

ΙT CD4-positive T cell

> (inhibition of pos. selection; preparation of aminoazepanones as inhibitors of Cathepsin L)

II

IT Antirheumatic agents

Rheumatoid arthritis

(preparation of aminoazepanones as inhibitors of Cathepsin L)

IT 350796-38-2P 350796-41-7.P 764650-55-7P RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES

```
(Uses)
        (drug candidate; preparation of aminoazepanones as inhibitors of Cathepsin
        L)
ΙT
     60616-82-2, Cathepsin L
     RL: BSU (Biological study, unclassified); BIOL (Biological study)
        (inhibition of; preparation of aminoazepanones as inhibitors of Cathepsin L)
ΙT
     150989-59-6P, 8-Oxa-3-azabicyclo[5.1.0]octane-3-carboxylic acid benzyl
             150989-61-0P, 4-Amino-3-hydroxyazepane-1-carboxylic acid benzyl
             150989-62-1P, 2,3,4,7-Tetrahydroazepine-1-carboxylic acid benzyl
             281219-32-7P
                           281219-33-8P, 4-Azido-3-hydroxyazepane-1-carboxylic
     ester
     acid benzyl ester
                         369593-24-8P
                                         369593-25-9P
                                                        764650-56-8P
     764650-57-9P
     RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
     (Reactant or reagent)
        (intermediate; preparation of aminoazepanones as inhibitors of Cathepsin L)
IT
     1119-51-3, 5-Bromo-1-pentene 5041-33-8 10349-57-2,
                                    13734-34-4
                                                               66715-65-9,
     Quinoline-6-carboxylic acid
                                                 58438-04-3
     Pyridin-2-sulfonyl chloride
     RL: RCT (Reactant); RACT (Reactant or reagent)
        (preparation of aminoazepanones as inhibitors of Cathepsin L)
     350796-38-2P
RN
RN
     350796-41-7P
RN
     764650-55-7P
     60616-82-2
RN
     150989-59-6P
RN
     150989-61-0P
RN
RN
     150989-62-1P
RN
     281219-32-7P
RN
     281219-33-8P
RN
     369593-24-8P
RN
     369593-25-9P
RN
     764650-56-8P
RN
     764650-57-9P
RN
     1119-51-3
RN
     5041-33-8
     10349-57-2
RN
RN
     13734-34-4
RN
     58438-04-3
RN
     66715-65-9
=> s 350796-38-2/rn or 350796-41-7/rn or 764650-55-7/rn
             4 350796-38-2
             0 350796-38-2D
             4 350796-38-2/RN
                 (350796-38-2 (NOTL) 350796-38-2D )
             4 350796-41-7
             0 350796-41-7D
             4 350796-41-7/RN
                 (350796-41-7 (NOTL) 350796-41-7D )
             1 764650-55-7
             0 764650-55-7D
             1 764650-55-7/RN
                 (764650-55-7 (NOTL) 764650-55-7D)
ь7
             4 350796-38-2/RN OR 350796-41-7/RN OR 764650-55-7/RN
=> dup rem 17
PROCESSING COMPLETED FOR L7
              4 DUP REM L7 (0 DUPLICATES REMOVED)
=> d ibib abs hitstr 1-4
rs
     ANSWER 1 OF 4 CAPLUS COPYRIGHT 2006 ACS on STN
```

ACCESSION NUMBER: 2004:803931 CAPLUS

DOCUMENT NUMBER: 141:295878

TITLE: Preparation of aminoazepanones as Cathepsin L

inhibitors

INVENTOR(S): Marquis, Robert W.

PATENT ASSIGNEE(S): USA

U.S. Pat. Appl. Publ., 13 pp. SOURCE:

CODEN: USXXCO

DOCUMENT TYPE:

Patent

LANGUAGE:

English

FAMILY ACC. NUM. COUNT:

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE		
US 2004192674	A1	20040930	US 2004-772817	20040205		
PRIORITY APPLN. INFO.:	ı		US 2003-447558P P	20030214		
OTHER SOURCE(S):	MARPAT	141:295878				
GI						

AΒ The title compds. I [R1 = substituted aminoalkylcarbonyl; R2 = H, alkyl, arylalkyl, etc.; R3 = H, alkyl, cycloalkyl, arylalkyl, etc.; R4 = H, alkyl, cycloalkyl, etc.] and their pharmaceutically acceptable salts are prepared and disclosed as inhibitors of Cathepsin L. Thus, e.g., II was prepared in a multistep synthesis employing N-Boc-phenylalanine. Consequently they are useful for preventing or treating diseases in which cathepsin L is implicated, such as rheumatoid arthritis or inhibition of pos. selection of CD4 + T-cells by cortical thymic epithelial cells.

IT 350796-38-2P 350796-41-7P 764650-55-7P

> RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(drug candidate; preparation of aminoazepanones as inhibitors of Cathepsin L)

ΙI

RN 350796-38-2 CAPLUS

CN 6-Quinolinecarboxamide, N-[(1S)-2-[[(4S)-hexahydro-3-oxo-1-(2pyridinylsulfonyl)-1H-azepin-4-yl]amino]-1-(2-naphthalenylmethyl)-2oxoethyl] - (9CI) (CA INDEX NAME)

RN 350796-41-7 CAPLUS

CN 6-Quinolinecarboxamide, N-[(1S)-2-[[(4S)-hexahydro-3-oxo-1-(2-pyridinylsulfonyl)-1H-azepin-4-yl]amino]-2-oxo-1-(phenylmethyl)ethyl]-(9CI) (CA INDEX NAME)

Absolute stereochemistry.

RN 764650-55-7 CAPLUS

CN 6-Quinolinecarboxamide, N-[(1S)-2-[[hexahydro-3-oxo-1-(2-pyridinylsulfonyl)-1H-azepin-4-yl]amino]-1-(2-naphthalenylmethyl)-2-oxoethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L8 ANSWER 2 OF 4 CAPLUS COPYRIGHT 2006 ACS on STN ACCESSION NUMBER: 2003:665404 CAPLUS

DOCUMENT NUMBER: 141:1037

TITLE: Potent and selective cathepsin L inhibitors do not

inhibit human osteoclast resorption in vitro. [Erratum

to document cited in CA135:120165]

AUTHOR(S): James, Ian E.; Marquis, Robert W.; Blake, Simon M.;

Hwang, Shing Mei; Gress, Catherine J.; Ru, Yu; Zembryki, Denise; Yamashita, Dennis S.; McQueney, Michael S.; Tomaszek, Thaddeus A.; Oh, Hye-Ja; Gowen,

Maxine; Veber, Daniel F.; Lark, Michael W.

CORPORATE SOURCE: Departments of Bone and Cartilage Biology, SmithKline

Beecham Pharmaceuticals, King of Prussia, PA, 19406,

USA

SOURCE: Journal of Biological Chemistry (2003), 278(34), 32484

CODEN: JBCHA3; ISSN: 0021-9258

PUBLISHER: American Society for Biochemistry and Molecular

Biology

DOCUMENT TYPE: Journal LANGUAGE: English

AB In Tables I and II, analogs SB-468430 and SB-468433 were originally reported to contain the quinoline-8-carboxamide moiety. Subsequent resynthesis revealed that in actuality these analogs contain the isomeric quinoline-6-carboxamide moiety. The modified structures are given in revised Tables I and II. Corrected Ki values for cathepsin L and cathepsin K in Table I are also given.

IT **350796-38-2**, SB 468430 **350796-41-7**, SB 468433

RL: BSU (Biological study, unclassified); BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(potent and selective cathepsin L inhibitors do not inhibit human

osteoclast resorption in vitro (Erratum))

RN 350796-38-2 CAPLUS

CN 6-Quinolinecarboxamide, N-[(1S)-2-[[(4S)-hexahydro-3-oxo-1-(2-pyridinylsulfonyl)-1H-azepin-4-yl]amino]-1-(2-naphthalenylmethyl)-2-oxoethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

RN 350796-41-7 CAPLUS

CN 6-Quinolinecarboxamide, N-[(1S)-2-[[(4S)-hexahydro-3-oxo-1-(2-pyridinylsulfonyl)-1H-azepin-4-yl]amino]-2-oxo-1-(phenylmethyl)ethyl]-(9CI) (CA INDEX NAME)

ANSWER 3 OF 4 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER:

2002:171694 CAPLUS

DOCUMENT NUMBER:

136:232208

TITLE:

Preparation of 4-aminoazepan-3-one parasitic cysteine protease inhibitors effective against malaria and

other diseases

CODEN: PIXXD2

INVENTOR(S):

Tew, David G.; Thompson, Scott K.; Veber, Daniel F.

Smithkline Beecham Corporation, UK

PATENT ASSIGNEE(S): SOURCE:

PCT Int. Appl., 220 pp.

DOCUMENT TYPE:

Patent

LANGUAGE:

English

FAMILY ACC. NUM. COUNT:

PATENT INFORMATION:

PA	PATENT NO.)	DATE	APPLICATION NO.					DATE				
WO	2002	0179	24		A1	-	2002	 0307_							2	00108	- 831
	W:	ΑE,	AG,	AL,	AM,	AT,	AU,	AZ,	BA,	BB,	BG,	BR,	BY,	ΒZ,	CA,	CH,	CN,
		co,	CR,	CU,	CZ,	DE,	DK,	DM,	DZ,	EC,	EE,	ES,	FI,	GB,	GD,	GE,	GH,
		GM,	HR,	HU,	ID,	IL,	IN,	IS,	JP,	ΚE,	KG,	KP,	KR,	ΚZ,	LC,	LK,	LR,
		LS,	LT,	LU,	LV,	MA,	MD,	MG,	MK,	MN,	MW,	MX,	MZ,	NO,	NZ,	PH,	PL,
		PT,	RO,	RU,	SD,	SE,	SG,	SI,	SK,	SL,	TJ,	TM,	TR,	TT,	TZ,	UA,	ŪG,
		US,	UZ,	VN,	YU,	ZA,	zw										
	RW:	GH,	GM,	KE,	LS,	MW,	ΜZ,	SD,	SL,	SZ,	TZ,	UG,	ZW,	ΑT,	BE,	CH,	CY,
		DE,	DK,	ES,	FI,	FR,	GB,	GR,	ΙE,	IT,	LU,	MC,	NL,	PT,	SE,	TR,	BF,
		ВJ,	CF,	CG,	CI,	CM,	GΑ,	GN,	GQ,	GW,	ML,	MR,	NE,	SN,	TD,	ΤG	
បន	2003	1441	75		A1	2003	0731	US 2001-881334					20010614				
AU	20010	08698	33		A5 20020313				AU 2001-86983					20010831			
EP	13203	370			Al		2003	0625	1	EP 2	001-9	9664	74		20	0108	331
	R:	ΑT,	BE,	CH,	DE,	DK,	ES,	FR,	GB,	GR,	IT,	LI,	LU,	NL,	SE,	MC,	PT,
		ΙE,	SI,	LT,	LV,	FI,	RO,	MK,	CY,	AL,	TR						
JP	20045	50908	33		Т2		2004	0325		JP 2	002-	52289	97		20	00108	331
PRIORIT'	Y APPI	LN. :	INFO.	. :					Ţ	JS 2	000-	6538:	15	I	A2 20	2000	901
									Ţ	JS 2	001-	88133	34	I	A2 20	00106	514
									Ţ	JS 1	998-	11363	36P	I	2 19	99812	223
									1	JS 1	999-:	16458	31P	I	2 19	9991	110
									1	VO 1	999-t	JS30	730	7	A2 19	99912	221
									Ţ	JS 2	000-	59384	45	I	32 20	00006	514
									Ţ	VO 2	00 1 -0	JS27:	178	V	V 20	0108	331
OTHER S	OURCE	(S):			MARI	PAT	136:2	23220	8(

GI

$$R^{1}$$
 R^{1}
 R^{1}
 R^{1}
 R^{2}
 R^{2}
 R^{4}
 R^{2}
 R^{4}
 R^{2}
 R^{4}
 R^{2}
 R^{4}
 R^{2}
 R^{4}
 R^{4}
 R^{4}
 R^{4}
 R^{4}
 R^{4}

AΒ The present invention relates to methods of treating parasitic diseases which are mediated by cysteine proteases by administration of 4-aminoazepan-3-one protease inhibitors I (e.g. benzo[1,3]dioxole-5carboxylic acid [(S)-1-(1-benzyl-3-oxoazepan-4-ylcarbamoyl)-3methylbutyl]amide) and pharmaceutically acceptable salts, hydrates and solvates thereof. In particular, the present invention relates to a method of treating malaria by inhibiting the cysteine protease falcipain. Other diseases against which the claimed compds. are effective include trypanosomiasis (African sleeping sickness, Chagas disease), leishmaniasis, schistosomiasis, onchocerciasis (river blindness) and giardiasis. In I: R1 is R4NR'CHR3C(O)-, R5XCHR3C(O)-, R3CH2C(O)-, R4NR'CR''''R3C(O)-, II. R2 is H, C1-6alkyl, C3-6cycloalkyl-C0-6alkyl, Ar-C0-6alkyl, Het-C0-6alkyl, R9C(0)-, R9C(S)-, R9SO2-, R9OC(0)-, R9R11NC(0), R9R11NC(S), R9(R11)NSO2, 3-(2-pyridyl)benzylcarbonyl, 2-(3-(2-pyridyl)phenyl)ethyl, R7NR6CHR8Z-, and R9SO2R11NC(O)-. R3 is H, C1-6alkyl, C3-6cycloalkyl-C0-6alkyl, C2-6alkenyl, C2-6alkynyl, HetCO-6alkyl and ArCO-6alkyl. R3 and R' may be connected to form a pyrrolidine, piperidine or morpholine ring. R4 is H, C1-6alkyl, C3-6cycloalkyl-C0-6alkyl, Ar-C0-6alkyl, Het-C0-6alkyl, R5C(0)-, R5C(S)-, R5SO2-, R5OC(0)-, R5R12NC(0)-, and R5R12NC(S)-. R5 is H, C1-6alkyl, C2-6alkenyl, C2-6alkynyl, C3-6cycloalkyl-C0-6alkyl, Ar-C0-6alkyl and Het-CO-6alkyl. R6 is H, C1-6alkyl, Ar-CO-6alkyl, and Het-CO-6alkyl. R7 is H, Cl-6alkyl, C3-6cycloalkyl-C0-6alkyl, Ar-C0-6alkyl, Het-C0-6alkyl, R10C(0)-, R10C(S)-, R10SO2-, R10OC(0)-, R10R13NC(0)-, and R10R13NC(S)-. R8 is H, C1-6alkyl, C2-6alkenyl, C2-6alkynyl, HetC0-6alkyl and Arco-6alkyl. R9, R10 independently = C1-6alkyl, C3-6cycloalkyl-C0-6alkyl, Ar-C0-6alkyl and Het-C0-6alkyl. R11, R12, R13, R', R'' independently = H, C1-6alkyl, Ar-C0-6alkyl, and Het-C0-6alkyl. R''' is H, C1-6alkyl, C3-6cycloalkyl-C0-6alkyl, Ar-C0-6alkyl, and Het-C0-6alkyl; R'''' is C1-6alkyl, C3-6cycloalkyl-C0-6alkyl C2-6alkenyl, C2-6alkynyl, HetC0-6alkyl and ArC0-6alkyl. X is CH2, S, and O; Z is C(O) and CH2; n is 1-5. Although the methods of preparation are not claimed, 220 example prepns. are included.

350796-38-2P, Quinoline-6-carboxylic acid [(1S)-2-(naphthalen-2-yl)-1-[[(4S)-3-oxo-1-(pyridin-2-ylsulfonyl)azepan-4-yl]carbamoyl]ethyl]amide 350796-41-7P, Quinoline-6-carboxylic acid [(1S)-1-[[(4S)-3-oxo-1-(pyridin-2-ylsulfonyl)azepan-4-yl]carbamoyl]-2-phenylethyl]amide
RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of 4-aminoazepan-3-one parasitic cysteine protease inhibitors effective against malaria and other diseases)

RN 350796-38-2 CAPLUS

CN

6-Quinolinecarboxamide, N-[(1S)-2-[[(4S)-hexahydro-3-oxo-1-(2-pyridinylsulfonyl)-1H-azepin-4-yl]amino]-1-(2-naphthalenylmethyl)-2-oxoethyl]- (9CI) (CA INDEX NAME)

RN 350796-41-7 CAPLUS

CN 6-Quinolinecarboxamide, N-[(1S)-2-[[(4S)-hexahydro-3-oxo-1-(2-pyridinylsulfonyl)-1H-azepin-4-yl]amino]-2-oxo-1-(phenylmethyl)ethyl]-(9CI) (CA INDEX NAME)

Absolute stereochemistry.

REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 4 OF 4 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2001:318582 CAPLUS

DOCUMENT NUMBER: 135:120165

TITLE: Potent and selective cathepsin L inhibitors do not

inhibit human osteoclast resorption in vitro

AUTHOR(S): James, Ian E.; Marquis, Robert W.; Blake, Simon M.;

Hwang, Shing Mei; Gress, Catherine J.; Ru, Yu; Zembryki, Denise; Yamashita, Dennis S.; McQueney, Michael S.; Tomaszek, Thaddeus A.; Oh, Hye-Ja; Gowen,

Maxine; Veber, Daniel F.; Lark, Michael W.

CORPORATE SOURCE: Departments of Bone and Cartilage Biology, SmithKline

Beecham Pharmaceuticals, King of Prussia, PA, 19406,

USA

SOURCE: Journal of Biological Chemistry (2001), 276(15),

11507-11511

CODEN: JBCHA3; ISSN: 0021-9258

PUBLISHER: American Society for Biochemistry and Molecular

Biology

DOCUMENT TYPE: Journal LANGUAGE: English

AB Cathepsins K and L are related cysteine proteases that have been proposed to play important roles in osteoclast-mediated bone resorption. To further examine the putative role of cathepsin L in bone resorption, we have evaluated selective and potent inhibitors of human cathepsin L and cathepsin K in an in vitro assay of human osteoclastic resorption and an

in situ assay of osteoclast cathepsin activity. The potent selective cathepsin L inhibitors (Ki = 0.0099, 0.034, and 0.27 nM) were inactive in both the in situ cytochem. assay (IC50 > 1 $\mu\text{M})$ and the osteoclast-mediated bone resorption assay (IC50 > 300 nM). Conversely, the cathepsin K selective inhibitor was potently active in both the cytochem. (IC50 = 63 nM) and resorption (IC50 = 71 nM) assays. A recently reported dipeptide aldehyde with activity against cathepsins L (Ki = 0.052 nM) and K (Ki = 1.57 nM) was also active in both assays (IC50 = 110 and 115 nM, resp.) These data confirm that cathepsin K and not cathepsin L is the major protease responsible for human osteoclastic bone resorption.

IT 350796-38-2 350796-41-7

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(potent and selective cathepsin L inhibitors do not inhibit human osteoclast resorption in vitro)

RN 350796-38-2 CAPLUS

CN 6-Quinolinecarboxamide, N-[(1S)-2-[[(4S)-hexahydro-3-oxo-1-(2-pyridinylsulfonyl)-1H-azepin-4-yl]amino]-1-(2-naphthalenylmethyl)-2-oxoethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

RN 350796-41-7 CAPLUS

CN 6-Quinolinecarboxamide, N-[(1S)-2-[[(4S)-hexahydro-3-oxo-1-(2-pyridinylsulfonyl)-1H-azepin-4-yl]amino]-2-oxo-1-(phenylmethyl)ethyl]-(9CI) (CA INDEX NAME)

Absolute stereochemistry.

REFERENCE COUNT:

21 THERE ARE 21 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

```
L1 ANSWER 1 OF 2 REGISTRY COPYRIGHT 2006 ACS on STN
```

RN 350796-41-7 REGISTRY

ED Entered STN: 09 Aug 2001

CN 6-Quinolinecarboxamide, N-[(1S)-2-[[(4S)-hexahydro-3-oxo-1-(2-pyridinylsulfonyl)-1H-azepin-4-yl]amino]-2-oxo-1-(phenylmethyl)ethyl]-(9CI) (CA INDEX NAME)

OTHER NAMES:

CN Quinoline-6-carboxylic acid [(1S)-1-[[(4S)-3-oxo-1-(pyridin-2-ylsulfonyl)azepan-4-yl]carbamoyl]-2-phenylethyl]amide

CN SB 468433

FS STEREOSEARCH

MF C30 H29 N5 O5 S

SR CA

LC STN Files: CA, CAPLUS, USPATFULL

Absolute stereochemistry.

PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

4 REFERENCES IN FILE CA (1907 TO DATE)

4 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L1 ANSWER 2 OF 2 REGISTRY COPYRIGHT 2006 ACS on STN

RN **350796-38-2** REGISTRY

ED Entered STN: 09 Aug 2001

CN 6-Quinolinecarboxamide, N-[(1S)-2-[[(4S)-hexahydro-3-oxo-1-(2-pyridinylsulfonyl)-1H-azepin-4-yl]amino]-1-(2-naphthalenylmethyl)-2-oxoethyl]- (9CI) (CA INDEX NAME)

OTHER NAMES:

CN Quinoline-6-carboxylic acid [(1S)-2-(naphthalen-2-yl)-1-[[(4S)-3-oxo-1-(pyridin-2-ylsulfonyl)azepan-4-yl]carbamoyl]ethyl]amide

CN SB 468430

FS STEREOSEARCH

MF C34 H31 N5 O5 S

SR CA

LC STN Files: CA, CAPLUS, USPATFULL

PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

- 4 REFERENCES IN FILE CA (1907 TO DATE) 4 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L4 ANSWER 1 OF 7 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2003:665404 CAPLUS

DOCUMENT NUMBER: 141:103796

TITLE: Potent and selective cathepsin L inhibitors do not

inhibit human osteoclast resorption in vitro. [Erratum

to document cited in CA135:120165]

AUTHOR(S): James, Ian E.; Marquis, Robert W.; Blake, Simon M.;

Hwang, Shing Mei; Gress, Catherine J.; Ru, Yu; Zembryki, Denise; Yamashita, Dennis S.; McQueney, Michael S.; Tomaszek, Thaddeus A.; Oh, Hye-Ja; Gowen,

Maxine; Veber, Daniel F.; Lark, Michael W.

CORPORATE SOURCE: Departments of Bone and Cartilage Biology, SmithKline

Beecham Pharmaceuticals, King of Prussia, PA, 19406,

USA

SOURCE: Journal of Biological Chemistry (2003), 278(34), 32484

CODEN: JBCHA3; ISSN: 0021-9258

PUBLISHER: American Society for Biochemistry and Molecular

Biology

DOCUMENT TYPE: Journal

LANGUAGE: English
AB In Tables I and II, analogs SB-468430 and SB

-468433 were originally reported to contain the quinoline-8-carboxamide moiety. Subsequent resynthesis revealed that in actuality these analogs contain the isomeric quinoline-6-carboxamide moiety. The modified structures are given in revised Tables I and II.

Corrected Ki values for cathepsin L and cathepsin K in Table I are also given.

IT 350796-38-2, SB 468430 350796-41-7,

SB 468433

RL: BSU (Biological study, unclassified); BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(potent and selective cathepsin L inhibitors do not inhibit human osteoclast resorption in vitro (Erratum))

RN 350796-38-2 CAPLUS

CN 6-Quinolinecarboxamide, N-[(1S)-2-[[(4S)-hexahydro-3-oxo-1-(2-pyridinylsulfonyl)-1H-azepin-4-yl]amino]-1-(2-naphthalenylmethyl)-2-oxoethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

RN 350796-41-7 CAPLUS

CN 6-Quinolinecarboxamide, N-[(1S)-2-[[(4S)-hexahydro-3-oxo-1-(2-pyridinylsulfonyl)-1H-azepin-4-yl]amino]-2-oxo-1-(phenylmethyl)ethyl]-(9CI) (CA INDEX NAME)

Absolute stereochemistry.

ANSWER 2 OF 7 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER:

2004:803931 CAPLUS

DOCUMENT NUMBER:

141:295878

TITLE:

Preparation of aminoazepanones as Cathepsin L

inhibitors

INVENTOR(S):

Marquis, Robert W.

PATENT ASSIGNEE(S):

USA:

SOURCE:

U.S. Pat. Appl. Publ., 13 pp.

CODEN: USXXCO

DOCUMENT TYPE:

Patent

LANGUAGE:

English

Ι

FAMILY ACC. NUM. COUNT:

PATENT INFORMATION:

PATENT NO.	KIND	DATE.	APPLICATION NO.	DATE
US 200419267		20040930	US 2004-772817	20040205
PRIORITY APPLN. I	NFO.:		US 2003-447558P P	20030214
OTHER SOURCE(S):	MARPAT	141:295878		
GI				

AΒ The title compds. I [R1 = substituted aminoalkylcarbonyl; R2 = H, alkyl, arylalkyl, etc.; R3 = H, alkyl, cycloalkyl, arylalkyl, etc.; R4 = H, alkyl, cycloalkyl, etc.] and their pharmaceutically acceptable salts are prepared and disclosed as inhibitors of Cathepsin L. Thus, e.g., II was prepared in a multistep synthesis employing N-Boc-phenylalanine. Consequently they are useful for preventing or treating diseases in which cathepsin L is implicated, such as rheumatoid arthritis or inhibition of

II

pos. selection of CD4 + T-cells by cortical thymic epithelial cells.

IT 350796-38-2P 350796-41-7P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(drug candidate; preparation of aminoazepanones as inhibitors of Cathepsin L)

RN 350796-38-2 CAPLUS


CN 6-Quinolinecarboxamide, N-[(1S)-2-[[(4S)-hexahydro-3-oxo-1-(2-pyridinylsulfonyl)-1H-azepin-4-yl]amino]-1-(2-naphthalenylmethyl)-2-oxoethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

RN 350796-41-7 CAPLUS

CN 6-Quinolinecarboxamide, N-[(1S)-2-[[(4S)-hexahydro-3-oxo-1-(2-pyridinylsulfonyl)-1H-azepin-4-yl]amino]-2-oxo-1-(phenylmethyl)ethyl]-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 3 OF 7 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2

2002:171694 CAPLUS

DOCUMENT NUMBER:

136:232208

TITLE:

Preparation of 4-aminoazepan-3-one parasitic cysteine protease inhibitors effective against malaria and

other diseases

INVENTOR(S):

Tew, David G.; Thompson, Scott K.; Veber, Daniel F.

PATENT ASSIGNEE(S): Smithkline Beecham Corporation, UK

SOURCE:

PCT Int. Appl., 220 pp.

DOCUMENT TYPE:

CODEN: PIXXD2

LANGUAGE:

Patent English

FAMILY ACC. NUM. COUNT:

111

PATENT INFORMATION:

PATENT NO.					KIN	D	DATE		APPLICATION NO.						DATE			
WO	2002	0179	24		A1		2002	0307			2001-				2	0010	831	
	W:	ΑE,	AG,	AL,	AM,	AT,	ΑU,	AZ,	BA,	BB	, BG,	BR,	BY,	ΒZ,	CA,	'CH,	CN,	
		CO,	CR,	CU,	CZ,	DE,	DK,	DM,	DZ,	EC	, EE,	ES,	FI,	GB,	GD,	GE,	GH,	
		GM,	HR,	HU,	ID,	ΙL,	IN,	IS,	JP,	KE	, KG,	ΚP,	KR,	ΚZ,	LC,	LK,	LR,	
		LS,	LT,	LU,	LV,	MA,	MD,	MG,	MK,	MN	, MW,	MX,	MZ,	NO,	NZ,	PH,	PL,	
		PT,	RO,	RU,	SD,	SE,	SG,	SI,	SK,	\mathtt{SL}	, TJ,	TM,	TR,	TT,	TZ,	UA,	UG,	
		US,	UZ,	VN,	YU,	ZA,	ZW											
	RW:	GH,	GM,	KE,	LS,	MW,	MZ,	SD,	SL,	SZ	, TZ,	UG,	ZW,	AT,	BE,	CH,	CY,	
		DE,	DK,	ES,	FI,	FR,	GB,	GR,	IE,	IT	, LU,	MC,	NL,	PT,	SE,	TR,	BF,	
		ВJ,	CF,	CG,	CI,	CM,	GΑ,	GN,	GQ,	GW	, ML,	MR,	NE,	SN,	TD,	ΤG		
US	US 2003144175				A1 20030731				US 2001-881334					20010614				
AU	2001										2001-							
EP	1320	370			A1		2003	0625		ΕP	2001-	9664	74		2	0010	831	
	R:	AT,	BE,	CH,	DE,	DK,	ES,	FR,	GB,	GR	, IT,	LI,	LU,	NL,	SE,	MC,	PT,	
		-	-		-	-					, TR							
						2004	0325	JP 2002-522897					20010831					
PRIORIT	Y APP	LN.	INFO	.:						US	2000-	6538	15	1	A2 2	0000	901	
											2001-			_	_	0010		
										US	1998-	1136	36P		P 1	9981	223	
											1999-					9991		
											1999-1			_				
											2000-							
										WO	2001-1	US27	178	1	W 2	0010	831	
OTHER S	OURCE	(S):			MAR	PAT	136:	23220	8 (

$$R^{1}$$
 R^{1}
 R^{1}
 R^{1}
 R^{2}
 R^{4}
 R^{2}
 R^{4}

The present invention relates to methods of treating parasitic diseases AΒ which are mediated by cysteine proteases by administration of 4-aminoazepan-3-one protease inhibitors I (e.g. benzo[1,3]dioxole-5carboxylic acid [(S)-1-(1-benzyl-3-oxoazepan-4-ylcarbamoyl)-3methylbutyl]amide) and pharmaceutically acceptable salts, hydrates and solvates thereof. In particular, the present invention relates to a method of treating malaria by inhibiting the cysteine protease falcipain. Other diseases against which the claimed compds. are effective include trypanosomiasis (African sleeping sickness, Chagas disease), leishmaniasis, schistosomiasis, onchocerciasis (river blindness) and giardiasis. In I: R1 is R4NR'CHR3C(0)-, R5XCHR3C(0)-, R3CH2C(0)-, R4NR'CR''''R3C(O)-, II. R2 is H, C1-6alkyl, C3-6cycloalkyl-C0-6alkyl, Ar-C0-6alkyl, Het-C0-6alkyl, R9C(0)-, R9C(S)-, R9SO2-, R9OC(O)-, R9R11NC(O)-, R9R11NC(S)-, R9(R11)NSO2-, 3-(2-pyridyl)benzylcarbonyl, 2-(3-(2-pyridyl)phenyl)ethyl, R7NR6CHR8Z-, and R9SO2R11NC(0)-. R3 is H, C1-6alkyl, C3-6cycloalkyl-C0-6alkyl, C2-6alkenyl, C2-6alkynyl, HetC0-6alkyl and ArC0-6alkyl. R3 and R' may be connected to form a

pyrrolidine, piperidine or morpholine ring. R4 is H, C1-6alkyl, C3-6cycloalkyl-C0-6alkyl, Ar-C0-6alkyl, Het-C0-6alkyl, R5C(0)-, R5C(S)-, R5SO2-, R5OC(O)-, R5R12NC(O)-, and R5R12NC(S)-. R5 is H, C1-6alkyl, C2-6alkenyl, C2-6alkynyl, C3-6cycloalkyl-C0-6alkyl, Ar-C0-6alkyl and Het-C0-6alkyl. R6 is H, C1-6alkyl, Ar-C0-6alkyl, and Het-C0-6alkyl. R7 is H, C1-6alkyl, C3-6cycloalkyl-C0-6alkyl, Ar-C0-6alkyl, Het-C0-6alkyl, R10C(O)-, R10C(S)-, R10SO2-, R10OC(O)-, R10R13NC(O)-, and R10R13NC(S)-. R8 is H, C1-6alkyl, C2-6alkenyl, C2-6alkynyl, HetC0-6alkyl and ArC0-6alkyl. R9, R10 independently = C1-6alkyl, C3-6cycloalkyl-C0-6alkyl, Ar-C0-6alkyl and Het-C0-6alkyl. R11, R12, R13, R', R'' independently = H, C1-6alkyl, Ar-C0-6alkyl, and Het-C0-6alkyl. R''' is H, C1-6alkyl, C3-6cycloalkyl-C0-6alkyl, Ar-C0-6alkyl, and Het-C0-6alkyl, R''' is C1-6alkyl, C3-6cycloalkyl-C0-6alkyl, C2-6alkenyl, C2-6alkynyl, HetC0-6alkyl and ArC0-6alkyl. X is CH2, S, and O; Z is C(O) and CH2; n is 1-5. Although the methods of preparation are not claimed, 220 example prepns. are included.

IT 350796-38-2P, Quinoline-6-carboxylic acid [(1S)-2-(naphthalen-2-yl)-1-[[(4S)-3-oxo-1-(pyridin-2-ylsulfonyl)azepan-4-yl]carbamoyl]ethyl]amide 350796-41-7P, Quinoline-6-carboxylic acid [(1S)-1-[[(4S)-3-oxo-1-(pyridin-2-ylsulfonyl)azepan-4-yl]carbamoyl]-2-phenylethyl]amide
RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of 4-aminoazepan-3-one parasitic cysteine protease inhibitors effective against malaria and other diseases)

·RN 350796-38-2 CAPLUS

CN

6-Quinolinecarboxamide, N-[(1S)-2-[[(4S)-hexahydro-3-oxo-1-(2-pyridinylsulfonyl)-1H-azepin-4-yl]amino]-1-(2-naphthalenylmethyl)-2-oxoethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

RN 350796-41-7 CAPLUS

CN 6-Quinolinecarboxamide, N-[(1S)-2-[[(4S)-hexahydro-3-oxo-1-(2-pyridinylsulfonyl)-1H-azepin-4-yl]amino]-2-oxo-1-(phenylmethyl)ethyl]-(9CI) (CA INDEX NAME)

RN 350796-41-7 CAPLUS

CN 6-Quinolinecarboxamide, N-[(1S)-2-[[(4S)-hexahydro-3-oxo-1-(2pyridinylsulfonyl)-1H-azepin-4-yl]amino]-2-oxo-1-(phenylmethyl)ethyl]-(9CI) (CA INDEX NAME)

Absolute stereochemistry.

REFERENCE COUNT:

2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 4 OF 7 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: DOCUMENT NUMBER:

2001:318582 CAPLUS

DOCUMENT

135:120165

TITLE:

Potent and selective cathepsin L inhibitors do not

inhibit human osteoclast resorption in vitro

AUTHOR(S):

James, Ian E.; Marquis, Robert W.; Blake, Simon M.; Hwang, Shing Mei; Gress, Catherine J.; Ru, Yu; Zembryki, Denise; Yamashita, Dennis S.; McQueney, Michael S.; Tomaszek, Thaddeus A.; Oh, Hye-Ja; Gowen,

Maxine; Veber, Daniel F.; Lark, Michael W.

CORPORATE SOURCE:

Departments of Bone and Cartilage Biology, SmithKline

Beecham Pharmaceuticals, King of Prussia, PA, 19406,

USA

SOURCE:

Journal of Biological Chemistry (2001), 276(15),

11507-11511

CODEN: JBCHA3; ISSN: 0021-9258

PUBLISHER:

American Society for Biochemistry and Molecular

Biology

DOCUMENT TYPE: LANGUAGE:

Journal English

AB Cathepsins K and L are related cysteine proteases that have been proposed to play important roles in osteoclast-mediated bone resorption. To further examine the putative role of cathepsin L in bone resorption, we have evaluated selective and potent inhibitors of human cathepsin L and cathepsin K in an in vitro assay of human osteoclastic resorption and an in situ assay of osteoclast cathepsin activity. The potent selective cathepsin L inhibitors (Ki = 0.0099, 0.034, and 0.27 nM) were inactive in both the in situ cytochem. assay (IC50 > 1 μM) and the osteoclast-mediated bone resorption assay (IC50 > 300 nM). Conversely, the cathepsin K selective inhibitor was potently active in both the cytochem. (IC50 = 63 nM) and resorption (IC50 = 71 nM) assays. A recently reported dipeptide aldehyde with activity against cathepsins L (Ki = 0.052) nM) and K (Ki = 1.57 nM) was also active in both assays (IC50 = 110 and 115 nM, resp.) These data confirm that cathepsin K and not cathepsin ${\tt L}$ is the major protease responsible for human osteoclastic bone resorption. IT Osteoclast

(potent and selective cathepsin L inhibitors do not inhibit human osteoclast resorption in vitro).

IT Bone

(resorption; potent and selective cathepsin L inhibitors do not inhibit human osteoclast resorption in vitro)

IT 60616-82-2, Cathepsin L 94716-09-3, Cathepsin K

RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(potent and selective cathepsin L inhibitors do not inhibit human osteoclast resorption in vitro)

IT 167498-29-5, SB 412515 251457-34-8, SB 290190 **350796-38-2** 350796-39-3 **350796-41-7**

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(potent and selective cathepsin L inhibitors do not inhibit human osteoclast resorption in vitro)

IT 350796-38-2 350796-41-7

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(potent and selective cathepsin L inhibitors do not inhibit human osteoclast resorption in vitro)

RN 350796-38-2 CAPLUS

CN 6-Quinolinecarboxamide, N-[(1S)-2-[[(4S)-hexahydro-3-oxo-1-(2-pyridinylsulfonyl)-1H-azepin-4-yl]amino]-1-(2-naphthalenylmethyl)-2-oxoethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

RN 350796-41-7 CAPLUS

CN 6-Quinolinecarboxamide, N-[(1S)-2-[[(4S)-hexahydro-3-oxo-1-(2-pyridinylsulfonyl)-1H-azepin-4-yl]amino]-2-oxo-1-(phenylmethyl)ethyl]-(9CI) (CA INDEX NAME)

Absolute stereochemistry.

REFERENCE COUNT:

21 THERE ARE 21 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 5 OF 7 EMBASE COPYRIGHT (c) 2006 Elsevier B.V. All rights

reserved on STN

ACCESSION NUMBER: 2003250069 EMBASE

TITLE: Osteoporosis: Challenges and new opportunities for therapy.

AUTHOR: Rotella D.P.

CORPORATE SOURCE: D.P. Rotella, Hopewell Discovery Chemistry, Bristol-Myers

Squibb Company, PO Box 5400, Princeton, NJ 08543 5400,

United States. david.rotella@bms.com

SOURCE: Current Opinion in Drug Discovery and Development, (2002)

Vol. 5, No. 4, pp. 477-486. .

Refs: 55

ISSN: 1367-6733 CODEN: CODDFF

COUNTRY: United Kingdom

DOCUMENT TYPE: Journal; General Review FILE SEGMENT: 030 Pharmacology

037 Drug Literature Index 038 Adverse Reactions Titles

LANGUAGE: English SUMMARY LANGUAGE: English

ENTRY DATE: Entered STN: 17 Jul 2003

Last Updated on STN: 17 Jul 2003

AB Osteoporosis is a chronic disease that affects a large number of both men and women and is characterized by a decrease in bone mass, as well as weakened bones. It causes a significant amount of morbidity and mortality in patients and is often only diagnosed after a fracture occurs. This review will highlight recent advances in the development of novel anabolic approaches for treatment of osteoporosis, such as parathyroid hormone (PTH), calcium sensing receptor modulators, statins and prostanoid receptor agonists. Selected antiresorptive targets (cathepsin K inhibitors and vitronectin receptor antagonists) will also be surveyed.

L4 ANSWER 6 OF 7 EMBASE COPYRIGHT (c) 2006 Elsevier B.V. All rights reserved on STN

ACCESSION NUMBER: 2002172171 EMBASE

TITLE: Thiol-dependent enzymes and their inhibitors: A review.

AUTHOR: Leung-Toung R.; Li W.; Tam T.F.; Karimian K.

CORPORATE SOURCE: R. Leung-Toung, Medicinal Chemistry Department, Apotex

Research Inc., 400 Ormont Drive, Toronto, Ont. M9L 1N9,

Canada. rleung@apotex.ca

SOURCE: Current Medicinal Chemistry, (2002) Vol. 9, No. 9, pp.

979-1002. . Refs: 190

ISSN: 0929-8673 CODEN: CMCHE7

COUNTRY: Netherlands

DOCUMENT TYPE: Journal; General Review

FILE SEGMENT: 029 Clinical Biochemistry

030 Pharmacology

037 Drug Literature Index

LANGUAGE: English SUMMARY LANGUAGE: English

ENTRY DATE: Entered STN: 30 May 2002

Last Updated on STN: 30 May 2002

AB Biological thiol-dependent enzymes have recently received extensive attention in the literature because of their involvement in a variety of physiopathological conditions. The active thiol groups of these enzymes are derived from the cysteine residues present. Hence, in a biological system, the selective reversible or irreversible inhibition of the activity of these enzymes by modification of the thiol moiety may potentially lead to the development of a chemotherapeutic treatment. Despite all the research efforts involved in the attempt to develop potential chemotherapeutic treatments for the major diseases involving cysteine proteases, there are in fact no such treatments available yet. However, AG7088 (1) an inhibitor of rhinovirus-3C is in phase II/III

Mar

clinical trial for the treatment of common cold and VX-740 (2, pralnacasan) an inhibitor of caspase-1 is in phase II clinical trial as an anti-inflammatory agent for rheumatoid arthritis. Several other cysteine protease inhibitors (i.e., cathepsin K, and S) are in pre-clinical evaluation or pre-clinical development. Structure-based drug design approaches have been instrumental in the development of these inhibitors. Intensive biochemical studies on the cysteine proteases have shed some light on some potential targets for therapeutic development. In addition, new techniques and new ideas are constantly emerging. As such, an up-to-date review of the literature on thiol-dependent enzymes as potential targets and their inhibitors designed from peptidic, modified peptidomimetic scaffolds and from small heterocyclic molecules is presented.

L4 ANSWER 7 OF 7 EMBASE COPYRIGHT (c) 2006 Elsevier B.V. All rights reserved on STN

ACCESSION NUMBER: 2002267120 EMBASE

TITLE: Recent advances in the synthesis, design and selection of

cysteine protease inhibitors.

AUTHOR: Alvarez Hernandez A.; Roush W.R.

CORPORATE SOURCE: A. Alvarez Hernandez, Department of Chemistry, University

of Michigan, Ann Arbor, MI 48109, United States.

roush@umich.edu

SOURCE: Current Opinion in Chemical Biology, (1 Aug 2002) Vol. 6,

No. 4, pp. 459-465. .

Refs: 46

ISSN: 1367-5931 CODEN: COCBF4

COUNTRY: United Kingdom

DOCUMENT TYPE: Journal; General Review FILE SEGMENT: 030 Pharmacology

037 Drug Literature Index

039 Pharmacy

LANGUAGE:

English

SUMMARY LANGUAGE:

=>

English

ENTRY DATE: Entered STN: 8 Aug 2002

Last Updated on STN: 8 Aug 2002

AB Inhibition of cysteine proteases is emerging as an important strategy for the treatment of a variety of human diseases. Intense efforts involving structure-based inhibitor design have been directed toward several cysteine proteases, including cathepsin K, calpain, human rhinovirus 3C protease and several parasitic cysteine protease targets. Other successful recent efforts have involved combinatorial synthesis and screening for identification of new inhibitor templates.